

### **BASIC CONTROL SYSTEMS**

02 LAPLACE TRANSFORM

**HANSHU YU** 

**NOVEMBER 2025** 



WHERE STUDENTS MATTER



## Presentation outline

Integral transform

Laplace transform

ROC of Laplace

Properties of Laplace transform



Inverse Laplace transform



#### **INTEGRAL TRANSFORM**

Why do we use integral transform?


Simplification

What does an integral transform do?

Mapping from one domain to another

Originates from:

Solving differential equations





#### **INTEGRAL TRANSFORM**

Given a general integral transform T

A function f(m) in m-domain

A target domain: *n*-domain

A transformation kernel: K(m, n)

The general integral transform:

$$\boldsymbol{L}[f(n)] = \int_{a}^{b} f(m)\boldsymbol{K}(m,n) \, \mathrm{d}m$$

We have mapped our function f from m domain to n domain using transformation T





### **LAPLACE TRANSFORM**

Laplace transform L

A function f(t) in t-domain

A target domain: s-domain

A transformation kernel:  $K(t,s) = e^{-st}$ 

Where  $s = \sigma + j\omega$ 

The Laplace transform:

$$\boldsymbol{L}[f(t)] = \int_{-\infty}^{\infty} f(t)e^{-st}dt$$



We have mapped our function f from t-domain to s-domain using  $\boldsymbol{L}$  .

This is call bilateral Laplace transform.



# UNILATERAL LAPLACE TRANSFORM

Recall that we are only working with causal system! So whatever before t = 0 we do not care, we start from 0!

The unilateral Laplace transform:

$$\boldsymbol{L}[f(t)] = \int_{0}^{\infty} f(t)e^{-st}dt$$

A transformation kernel:  $K(t,s) = e^{-st}$ Where  $s = \sigma + j\omega$ 





# UNILATERAL LAPLACE TRANSFORM

The unilateral Laplace transform:

$$\boldsymbol{L}[f(t)] = \int_{0}^{\infty} f(t)e^{-st}dt$$

A transformation kernel:  $K(t,s) = e^{-st}$ Where  $s = \sigma + j\omega$ 

**BUT WHY???** 





# CONVERGE! CONVERGE!

We are integrating from 0 to  $\infty$ . So f(t)K(s,t) must converge for out integral to exist.

The unilateral Laplace transform:

$$\boldsymbol{L}[f(t)] = \int_{0}^{\infty} f(t)e^{-st}dt$$

A transformation kernel:  $K(t,s) = e^{-st}$ Where  $s = \sigma + j\omega$ 





The unilateral Laplace transform:

$$L[f(t)] = \int_{0}^{\infty} f(t)e^{-st}dt$$
 We are integrating from 0 to  $\infty$ .

A transformation kernel:

$$K(t,s) = e^{-st}$$
  
Where  $s = \sigma + j\omega$ 

So f(t)K(s,t) must converge for out integral to exist.

We look at f(t) and K(s, t)separately, K(s, t) first.

Assume: 
$$f(t) = 1$$

$$\int_{0}^{\infty} e^{-st} dt = \int_{0}^{\infty} e^{-(\sigma+j\omega)t} dt = \int_{0}^{\infty} e^{-\sigma t} e^{-j\omega t} dt$$

Remember from complex analysis:

$$|e^{-j\omega t}| = 1$$

Then we only have:

$$e^{-\sigma t}$$



To converge:  $\sigma > 0$ 



The unilateral Laplace transform:

$$\boldsymbol{L}[f(t)] = \int_{0}^{\infty} f(t)e^{-st}dt$$

A transformation kernel: Assume: s = 0 + j0

$$K(t,s)=e^{-st}$$

Where  $s = \sigma + j\omega$ 

$$\int_{0}^{\infty} f(t) dt$$

Without our kernel, f(t) should converge.






The unilateral Laplace transform:

$$\boldsymbol{L}[f(t)] = \int_{0}^{\infty} f(t)e^{-st}dt$$

Where  $s = \sigma + j\omega$ 

A transformation kernel: Assume:  $s = 0 + j\omega$  $K(t,s) = e^{-st}$ 



We are traversing through the complex plane now, but our ROC still relies on the properties of f(t).





The unilateral Laplace transform:

$$\boldsymbol{L}[f(t)] = \int_{0}^{\infty} f(t)e^{-st} dt$$

A transformation kernel:

$$K(t,s) = e^{-st}$$

 $K(t,s) = e^{-st}$  Assume:  $s = \sigma + j\omega$ 

Where  $s = \sigma + j\omega$ 

$$\int_{0}^{\infty} e^{-\sigma t} f(t)e^{-j\omega t} dt \text{ , and } |e^{-j\omega t}| = 1$$

Finally, when  $\sigma > 0$ , we have an "envelop"  $e^{-\sigma t}$  that make our transformation exist.

This gives us the desired properties of f(t)

1. f(t) should be integrable and defined for  $[0, \infty)$ 

2. f(t) grows slower than  $e^{-st}$ 



# REGION OF CONVERGENCE THE "POLAR VIEW"

For any real number  $r = a\vec{x} + b\vec{y}$ , the polar form:  $r = |r| \angle r$ 

For any complex number z = a + jb, the polar form:

$$\mathbf{z} = |z|\hat{z}, \ \hat{z} = e^{j\theta}, \theta = \text{Arg}(z)$$

For our  $f(t)\mathbf{K}(s,t) = f(t)e^{-(\sigma+j\omega)t}$ :

We can obtain a polar representation:


$$|z|e^{i\theta}$$

$$|z| = e^{-\sigma t}f(t)$$

$$\theta = -\omega t$$

Putting them together:

$$f(t)\mathbf{K}(s,t) = [e^{-\sigma t}f(t)]e^{j(-\omega t)}$$






## **KERNEL OF LAPLACE**

$$\mathbf{K}(s,t) = f(t)e^{-(\sigma+\mathrm{j}\omega)t}$$

This is a very beautiful **probing function** for measuring our



This probe gives us very nice analytical properties such that we can map many time domain signals into complex space.

In the meantime, this probe still carries enough information such that we can recover the real signal from complex plane. (*Lerch's theorem*)



Laplace transform is **ONE-TO-ONE** (injective).



# CONVERGE! CONVERGE!

Impatient guy from engineering department rushed to me:

"THESE ARE JUST MATHEMATICAL TRICKS! Mathematical rules and all! Does not make sense in practice!"

"How does these relate to application & real systems?"

"WHY should we care ????"

I say:



"Because of the physical meanings"



# RECALL: SYSTEMS: THE KEY TAKEOUT

No matter how much we simplify, we are working with physical systems.

The mathematical tools you see later, are describing the characteristics of the physical system.

(最重要的是物理系统自身的特性!)





# ACTUAL PHYSICAL SYSTEMS OF INFINITE ENERGY DOES NOT EXIST!

**Energy = Power over TIME.** 

$$\mathbf{E} = \int_{0}^{\infty} \mathbf{P} \, \mathrm{d}\mathbf{t}$$

As t->  $\infty$ , diverging power, infinite energy, impossible. If this is the governing equation, the system will break!

**Unstable!** 





#### **EXAMPLE**

$$f(t) = e^{at}$$

Laplace transform:

$$\mathbf{F}(s) = \mathbf{L}[f(t)] = \int_{0}^{\infty} f(t)e^{-st} dt$$

$$= \int_{0}^{\infty} e^{at}e^{-st} dt$$

$$= \int_{0}^{\infty} e^{(a-s)t} dt$$

Without getting the exact F(s), we can already infer where is the ROC:



$$a - \sigma < 0$$

Thus:



#### **EXAMPLE**

$$f(t) = e^{at}$$

Laplace transform:

$$\mathbf{F}(s) = \mathbf{L}[f(t)] = \int_{0}^{\infty} f(t)e^{-st} dt$$

$$= \int_{0}^{\infty} e^{at} e^{-st} dt$$

$$= \int_{0}^{\infty} e^{(a-s)t} dt$$


$$a - \sigma < 0$$

$$\sigma > a$$



Thus:

The ROC:





#### **EXAMPLE**

$$f(t) = e^{at}$$

Laplace transform:

$$\mathbf{F}(s) = \mathbf{L}[f(t)] = \int_{0}^{\infty} f(t)e^{-st} dt$$

$$= \int_{0}^{\infty} e^{(a-s)t} dt$$

$$= \frac{1}{a-s} e^{(a-s)t} \Big|_{t=0}^{\infty}$$

$$= 0 - \frac{1}{a-s}$$

$$= \frac{1}{a-s} e^{(a-s)t} \Big|_{t=0}^{\infty}$$





#### LAPLACE TRANSFORM TABLE

1. 
$$A \cdot l(t)$$

2. 
$$\delta(t)\cdot I(t)$$

3. 
$$t^n \cdot I(t)$$

**4.** 
$$e^{at} \cdot I(t)$$

5. 
$$sin(\omega t) \cdot I(t)$$

**6.** 
$$cos(\omega t) \cdot I(t)$$

$$\frac{A}{s}$$

$$\frac{n!}{s^{n+1}}$$

$$\frac{I}{s-a}$$

$$\frac{\omega}{s^2 + \omega^2}$$

$$\frac{s}{s^2 + \omega^2}$$





## LAPLACE TRANSFORM TABLE

| f(t)                      | $F(s) = \mathcal{L}[f(t)]$           |          |
|---------------------------|--------------------------------------|----------|
| f(t) = 1                  | $F(s) = rac{1}{s}$                  | s > 0    |
| $f(t) = e^{at}$           | $F(s) = \frac{1}{(s-a)}$             | s > a    |
| $f(t)=t^n$                | $F(s) = \frac{n!}{s^{(n+1)}}$        | s > 0    |
| $f(t)=\sin(at)$           | $F(s) = \frac{a}{s^2 + a^2}$         | s > 0    |
| $f(t) = \cos(at)$         | $F(s) = \frac{s}{s^2 + a^2}$         | s > 0    |
| $f(t)=\sinh(at)$          | $F(s) = \frac{a}{s^2 - a^2}$         | s >  a   |
| $f(t) = \cosh(at)$        | $F(s) = \frac{s}{s^2 - a^2}$         | s >  a   |
| $f(t)=t^ne^{at}$          | $F(s) = \frac{n!}{(s-a)^{(n+1)}}$    | s > a    |
| $f(t) = e^{at}\sin(bt)$   | $F(s) = \frac{b}{(s-a)^2 + b^2}$     | s > a    |
| $f(t) = e^{at}\cos(bt)$   | $F(s) = \frac{(s-a)}{(s-a)^2 + b^2}$ | s > a    |
| $f(t)=e^{at}\sinh(bt)$    | $F(s) = \frac{b}{(s-a)^2 - b^2}$     | s-a >  b |
| $f(t) = e^{at} \cosh(bt)$ | $F(s) = \frac{(s-a)}{(s-a)^2 - b^2}$ | s-a >  b |





#### **INVERSE LAPLACE TRANSFORM**

The time domain signal can be obtained from the frequency domain signal using inverse Laplace transform.

$$f(t) = L^{-1} [F(s)] = \frac{1}{2\pi j} \int_{\sigma - j\infty}^{\sigma + j\infty} F(s)e^{st} ds$$

Without transform table, you can use *Cauchy's residue theorem* for a contour integral with the closed contour of the integration as the region of convergence.



Luckily you will have the table in the formula sheet in the exam.



$$\boldsymbol{L}[f(t)] = \int_{0}^{\infty} f(t)e^{-st}dt$$

#### LINEARITY

$$a \cdot f(t) \Leftrightarrow a \cdot F(s)$$

$$f(t) + g(t) \Leftrightarrow F(s) + G(s)$$

$$a \cdot f(t) + b \cdot g(t) \Leftrightarrow a \cdot F(s) + b \cdot G(s)$$






$$\boldsymbol{L}[f(t)] = \int_{0}^{\infty} f(t)e^{-st} dt$$

TIME SCALING

$$f(at) \Leftrightarrow \frac{1}{|a|} \cdot F\left(\frac{s}{a}\right)$$

TIME SHIFTING

$$f(t-a) \Leftrightarrow e^{-as} \cdot F(s)$$








## Damping (example)

$$f(t) = e^{-at}cos(\omega t)$$



What do we need to find the laplace transform of f(t)?

Laplace transform theorems:

$$e^{-at}f(t)$$

$$F(s+a)$$

Laplace transform table:

**4.** 
$$e^{at} \cdot I(t)$$

$$\frac{I}{s-a}$$

**6.** 
$$cos(\omega t) \cdot l(t)$$

$$\frac{s}{s^2+\omega^2}$$



$$\mathcal{L}\lbrace e^{-at}\cos(\omega t)\rbrace = \frac{(s+a)}{(s+a)^2 + \omega^2}$$



$$\boldsymbol{L}[f(t)] = \int_{0}^{\infty} f(t)e^{-st}dt$$

#### DIFFERENTATION

$$L[f'(t)] = sF(s) - f(0)$$
  

$$L[f''(t)] = s^2F(s) - sf(0) - f'(0)$$

General formula:

$$L[f^{(k)}(t)] = s^k F(s) - s^{k-1} f(0) - s^{k-2} f'^{(0)} \dots - s f^{(k-2)}(0) - f^{(k-1)}(0)$$

Differentiation in t-domain in becomes an operator in s-domain.





$$\boldsymbol{L}[f(t)] = \int_{0}^{\infty} f(t)e^{-st}dt$$

#### **INTEGRAL**

$$g(t) = \int_0^t f(\tau) d\tau$$

With Laplace transform:

$$G(s) = \frac{1}{s} F(s)$$

Integration in t-domain in becomes an operator in s-domain.



$$\boldsymbol{L}[f(t)] = \int_{0}^{\infty} f(t)e^{-st}dt$$

#### From CONVOLUTION to MULTIPLICATION

$$f(t) * g(t) = \int_0^t f(\tau)g(t-\tau) d\tau$$

With Laplace transform:

$$f(t) * g(t) \Leftrightarrow F(s)G(s)$$

Convolution in t-domain in becomes multiplication in s-domain.





$$\boldsymbol{L}[f(t)] = \int_{0}^{\infty} f(t)e^{-st}dt$$

#### INITIAL VALUE THEOREM

$$\lim_{t\to 0} f(t) = \lim_{s\to \infty} sF(s)$$

Proof1:

f is causal & bounded such that  $\lim_{t\to 0^+} f(t) \to \alpha$ , we play with definition

$$F(s) = \int_{0}^{\infty} f(t)e^{-st} dt = \int_{0}^{\infty} f\left(\frac{t}{s}\right)e^{-s\frac{t}{s}} d\frac{t}{s} = \int_{0}^{\infty} \frac{1}{s} f\left(\frac{t}{s}\right)e^{-t} dt$$
$$sF(s) = \int_{0}^{\infty} f\left(\frac{t}{s}\right)e^{-t} dt$$

Based on Lebesgue's dominated convergence theorem, we can have

$$\lim_{s \to \infty} sF(s) = \int_{0}^{\infty} \alpha e^{-t} dt = \alpha = \lim_{t \to 0^{+}} f(t)$$



$$\boldsymbol{L}[f(t)] = \int_{0}^{\infty} f(t)e^{-st}dt$$

#### **INITIAL VALUE THEOREM**


$$\lim_{t\to 0} f(t) = \lim_{s\to \infty} sF(s)$$

Proof2:

f is causal & bounded such that  $\lim_{t\to 0^+} f(t) \to \alpha$ , we play with definition

$$F(s) = \int_{0}^{\infty} f(t)e^{-st} dt = \int_{0}^{\infty} f\left(\frac{t}{s}\right)e^{-s\frac{t}{s}} d\frac{t}{s} = \int_{0}^{\infty} \frac{1}{s} f\left(\frac{t}{s}\right)e^{-t} dt$$
$$sF(s) = \int_{0}^{\infty} f\left(\frac{t}{s}\right)e^{-t} dt$$

We select a  $\delta \in \mathbb{R}$  sufficiently close to 0 such that  $\int_{\delta}^{\infty} e^{-t} dt < \epsilon$  that is arbitrarily small and  $\lim_{s\to\infty} f\left(\frac{t}{s}\right) = \alpha$  for  $t\in(0,\delta]$ , we may also conclude that:







$$\boldsymbol{L}[f(t)] = \int_{0}^{\infty} f(t)e^{-st}dt$$

#### **FINAL VALUE THEOREM**

$$\lim_{t \to \infty} f(t) = \lim_{s \to 0} sF(s)$$

Proof:

f is continuously differentiable and bounded, and f is absolutely integrable.

 $\lim_{t\to\infty} f(t)$  exists and finite.

$$\lim_{s \to 0} sF(s) = \lim_{s \to 0} L\left[\frac{\mathrm{d}f(t)}{\mathrm{d}t}\right] = \lim_{s \to 0} \int_{0}^{\infty} \frac{\mathrm{d}f(t)}{\mathrm{d}t} e^{-st} \mathrm{d}t$$

As obviously  $\lim_{s\to 0} e^{-st} = 1$ , we have:



$$\lim_{s \to 0} \int_{0}^{\infty} \frac{\mathrm{d}f(t)}{\mathrm{d}t} e^{-st} \mathrm{d}t = f(t) \Big|_{0}^{\infty} = f(\infty) - f(0)$$



We have seen the incredible symmetry between time domain and the frequency domain.

- Differentiation in one becomes multiplication in another.
- Exponential scaling in one domain becomes shifting in the other.





#### WARNING

We do not allow the following expressions to appear:

Although Fourier transform and Laplace transform looks similar, whoever says that:

Fourier = Laplace when  $s=j\omega$  or Laplace = Fourier when  $s=\sigma+j\omega$ 

Is absolutely WRONG!!!





#### WARNING

We do not allow the following expressions to appear:

Although Fourier transform and Laplace transform looks similar, whoever says that:

Fourier = Laplace when 
$$s=j\omega$$
  
Laplace = Fourier when  $s=\sigma+j\omega$ 


Is absolutely WRONG!!!

(But in practice this usually works.)





#### Modelling example: RC low-pass filter



Differential Equation! -> Laplace transform!


$$U_{in}(t) = RC \frac{dU_{out}}{dt} + U_{out}(t)$$
  

$$U_{in}(s) = RCsU_{out} + U_{out}(s)$$



$$U_{in} = RCsU_{out} + U_{out}$$





Laplace transform! -> Transfer function!

$$OUTPUT = INPUT \cdot TF$$


$$\Rightarrow$$

$$TF = \frac{OUTPUT}{INPUT}$$



$$H(s) = \frac{1}{1 + RCs}$$





Time domain solution?

Inverse Laplace transform!

$$H(s) = \frac{1}{1 + RCs}$$


$$U_{out}(s) = H(s)U_{in}(s)$$

We assume  $U_{in} = \frac{1}{s}$  (Constant)



$$U_{out}(s) = \frac{1}{s(1 + RCs)}$$





Inverse Laplace transform!


$$U_{out}(s) = \frac{1}{s(1 + RCs)}$$

Partial fraction decomposition:

$$U_{out}(s) = \frac{A}{s} + \frac{B}{1 + RCs}$$
$$A + (ARC + B)s = 1$$
$$A = 1, B = -RC$$








$$U_{out}(s) = \frac{1}{s} - \frac{RC}{1 + RCs} = \frac{1}{s} - \frac{1}{\frac{1}{RC} + s}$$





$$U_{out}(t) = 1 - e^{-\frac{t}{RC}}$$





### Circuit analysis! "The smart way"

| Time - domain   | S-domain                              |
|-----------------|---------------------------------------|
| Integration     | Multiplicative operator $\frac{1}{s}$ |
| Differentiation | Multiplicative operator 's'           |






# **ELECTRICAL SYSTEMS – RLC CIRCUITS**

|           | Voltage - Current                           | Impedance<br>(Laplace transformed) |
|-----------|---|------------------------------------|
| Resistor  | U(t) = I(t)R                                | R                                  |
| Capacitor | $U(t) = \frac{1}{C} \int_0^1 I(\tau) d\tau$ | $\frac{1}{Cs}$                     |
| Inductor  | $U(t) = L \frac{d I(t)}{d t}$               | Ls                                 |








### Circuit analysis! "The smart way"

|          | Resistive components | S-domain Impedance |
|----------|----------------------|--------------------|
|          | Resistor             | R                  |
|          | Inductor             | Ls                 |
| <b>Q</b> | Capacitor            | $\frac{1}{Cs}$     |







Circuit analysis!

Total resistance:  $R + \frac{1}{cs}$  over  $U_{in}$ 

 $U_{out}$  is the voltage over resistance  $\frac{1}{Cs}$ .

Like a simple pure resistor circuit:



$$\frac{\mathsf{U}_{\mathsf{out}}}{\mathsf{U}_{\mathsf{in}}} = \frac{\frac{1}{CS}}{R + \frac{1}{CS}} = \frac{1}{1 + RCS}$$




## Can you create a low-pass filter with a resistor and an inductor?

Try it yourself!





### Modelling example: Mechanical system



Write differential equation

$$F(t) = k \int_{0}^{t} v(\tau) d\tau + bv(t) + m \frac{dv(t)}{dt}$$


Laplace transform

$$F(s) = \frac{k}{s}V(s) + bV(s) + msV(s)$$





### Modelling example: Mechanical system



$$F(s) = \frac{k}{s}V(s) + bV(s) + msV(s)$$

Transfer function

$$\frac{V(s)}{F(s)} = \frac{1}{\frac{k}{s} + b + ms}$$
$$= \frac{s}{ms^2 + bs + k}$$





# MECHANICAL SYSTEMS – TRANSLATING SYSTEM

|                              | Force - Velocity                           | Impedance<br>(Laplace transformed) |
|------------------------------|--|------------------------------------|
| Damper<br>(Viscous friction) | F = bv                                     | b                                  |
| Spring                       | $F = k \int_{0}^{t} v(\tau) d\tau$         | $\frac{k}{s}$                      |
| Mass (Inertia)               | $F = m \frac{\mathrm{d}v(t)}{\mathrm{d}t}$ | ms                                 |





# MECHANICAL SYSTEMS – TRANSLATING SYSTEM

|                              | Force – Displacement                            | Impedance<br>(Laplace transformed) |
|------------------------------|---|------------------------------------|
| Damper<br>(Viscous friction) | $F = b \frac{\mathrm{d}x(t)}{\mathrm{d}t}$      | bs                                 |
| Spring                       | F = kx  | k                                  |
| Mass (Inertia)               | $F = m \frac{\mathrm{d}^2 x(t)}{\mathrm{d}t^2}$ | $ms^2$                             |





### **SUMMARY**

$$\boldsymbol{L}[f(t)] = \int_{0}^{\infty} f(t)e^{-st} dt$$

**Transform table** 

Region of convergence

### **Properties:**

- Linearity
- Symmetry between t-domain and s-domain
- From differentiation and integral to operators in sdomain
- Initial & final value theorem



We may use Laplace transform to conveniently solve ODE





## **SUMMARY**

The physics laws governs our systems' performance

Analyze your system based on these physics laws

Eventually we can use ODE to describe our desired input and output

Laplace transform is a handy tool for analysis and finding solutions





## **HOMEWORK**

Read appendix A

Knowledge Recap: Complex Analysis

You may skip the following contents:

- Italic contents on page 22
- Section A.D Why complex numbers?

They are fun but not essential to our course. But still, for those interested, it is highly recommended that you take a look.

#### Stage ONE exercise:

- Problem 4 (you can skip subquestion 4 and 5 for now, 5 is possible to do, so give it a try)
- Problem 5
- Problem 6
- Problem 7

